

ALPINE FLORA OF CERRO MOHINORA, CHIHUAHUA, MEXICO

J. Andrew McDonald

*Department of Biology
The University of Texas – Pan American
1201 W. University Dr.
Edinburg, Texas 78739, U.S.A.
amcdonald@utpa.edu*

Jorge Martinez

*Departamento de Biología
Centro de Ciencias Básicas, Edificio 202
Avenida Universidad #940
Ciudad Universitaria Aguascalientes
Aguascalientes, MÉXICO C.P. 20131*

Guy L. Nesom

*2925 Hartwood Drive
Fort Worth, Texas 76109, U.S.A.*

ABSTRACT

An annotated, vouchered checklist for the only known alpine-subalpine plant community in the Sierra Madre Occidental of Mexico, Cerro Mohinora of southwest Chihuahua, is presented. Contrary to former reports of a sparse timberline vegetation on Cerro Mohinora, we document a relatively diverse vascular flora, including 79 species that represent 63 genera and 33 families. Based on the species composition of this flora, the vegetation shares minimal biogeographic affinities with alpine vegetation in the Sierra Madre Oriental of Northeast Mexico (7.2% Sorensen Index; comparing Mohinora with the collective floras of Sierra La Viga, Coahuilón, Cerro Potosí, and Peña Nevada). Only nine timberline species of Mohinora occur in similar habitats in the high mountains of northeast Mexico, suggesting enduring isolation through time, even during pluvial periods of the Pleistocene. Cerro Mohinora shares about half that index of similarity with alpine vegetation in Mexico's volcanic belt (3.6% Sorensen Index; comparing Mohinora with the collective floras of Popocatepetl, Iztaccihuatl and Nevado de Toluca).

RESUMEN

Se presenta un listado florístico basado en colecciones vegetales para el único refugio alpino conocido en la Sierra Madre Occidental de México, Cerro Mohinora, en el suroeste de Chihuahua. A pesar de reportes históricos que describen una flora con pocas especies en este sitio, reportamos una comunidad vegetal que soporta una flora diversa con 79 especies, 63 géneros, y 33 familias. Basada en la composición de especies de esta flora, la vegetación alpina de Cerro Mohinora demuestra afinidades fitogeográficas mínimas con la vegetación alpina de la Sierra Madre Oriental. Se calcula un índice de similitud de 7.2% (Sorensen) entre la flora de Mohinora y las de Sierra La Viga, Coahuilón, Cerro Potosí, y Peña Nevada. Solamente nueve especies alpinas de Cerro Mohinora se encuentran en las montañas altas del Nordeste de México, indicando un aislamiento permanente, aun durante ciclos pluviales durante el Pleistoceno. La zona alpina de Cerro Mohinora demuestra aun menos afinidades con floras similares en la zona neovolcánica, demostrando un índice de similitud (Sorensen) de 3.6% entre la zona alpina de Mohinora y los zacatonales de Popocatepetl, Iztaccihuatl y el Nevado de Toluca.

INTRODUCTION

Studies on the alpine floras of Mexico focus primarily on highly specialized timberline elements that inhabit basaltic substrates above 4000 m on the 19° latitude parallel (Zone III, Fig. 1), including those of the Cofre de Perote, Iztaccihuatl, Malinche, Nevado de Colima, Nevado de Toluca, Pico de Orizaba, and Popocatepetl (Purpus 1907; Beaman 1962, 1965; Lauer & Klaus 1975; Narave 1985; Almeida-Leñero et al. 1994, 2004; Gimenez de Azcarate & Escamilla 1999). Beaman & Andresen (1966) expanded the scope of alpine studies by comparing the floristic composition of these volcanic peaks with an isolated community of alpine plants in Northeast Mexico, where timberline communities of Cerro Potosí occur on limestone substrates in the Sierra Madre Oriental of Nuevo Leon. This northern site exhibits closer phytogeographic affinities to forb-dominated montane plant communities of the southern Rocky Mountains than with grass-dominated, timberline vegetation ('zacatonales') across Mexico's volcanic belt (Beaman 1965; Rzedowski 1975; Narave 1985; Almeida-Leñero et al. 2004). Later explorations of Northeast Mexico by McDonald (1990, 1993) discovered six additional alpine sites in the region (three sites indicated in Fig. 1 on account of the close proximity of four timberline zones), between which a considerable number of autochthonous (McDonald 1993, Table 23.1), narrow endemic species occur.

None of the aforementioned studies make reference to the reconnaissance work of Donovan Correll (1960) and Howard Scott Gentry on an isolated peak of the Sierra Madre Occidental in Southwest Chihuahua (Fig. 1), Cerro Mohinora (25°57'29.31"N, 107°02'56.37"W), where Goldman (1951) had observed in 1898 a




FIG. 1. Contemporary distribution of alpine-subalpine floras in Mexico. A single, relictual timberline vegetation occurs in the Sierra Madre Occidental (Zone I: Cerro Mohinora), as compared to a series of closely situated alpine refugia in the Sierra Madre Oriental (Zone II). The central volcanic belt traversing the 19° lat. meridian (Zone III) and an isolated, highly restricted alpine/subalpine site on Cerro Quiexobra of the Sierra Madre del Sur (Zone IV, in prep harbor isolated and uniquely derived timberline vegetation).

snow-capped summit with stunted trees and 'open' forest. Arriving on the 16th of October, 1959, Correll reported a "disappointingly sparse" vegetation dominated by herbs and grasses on Cerro Mohinora's precipitous, rocky substrates (maximum altitude reports ranging from 3319–3900 m; Webb & Baker 1984). He attributed the peak's poor floristic composition to either goat grazing or a heavy frost that had already fallen by the time of his late arrival.

In recognition that the timberline vegetation of Cerro Mohinora had yet to be observed during the peak of flowering in late summer, the site was visited during the last two weeks of August in 1987 and 1988 by the authors to assess the floristic composition of the Sierra Madre Occidental's only known timberline flora. We observed a colorful and species-rich alpine vegetation on approximately 15 hectares upon the mountain's bald and adjacent declivities. Forbs and grasses dominate the timberline vegetation, representing 33 families, 63 genera, and 79 species, with an occasional intrusion of stunted pines (*Pinus cooperi* Blanco), junipers (*Juniperus blancoi* Martinez var. *huehuetentensis* R.P. Adams) and aspen (*Populus tremuloides* Michx.). The Asteraceae and Poaceae harbor more species than other vascular plant families (14 and 7 spp., respectively), followed by the Caryophyllaceae (4 spp.), Plantaginaceae (4 spp.), Apiaceae (3 spp.), Crassulaceae (3 spp.) and Primulaceae (3 spp.).

While the timberline floras of the Sierra Madre Oriental and Sierra Madre Occidental cover no more than

6 sq. kms in the present day, McDonald (1993) estimated approximately 5% of Mexico was covered by alpine vegetation during pluvial cycles of the Pleistocene. Based on the hypothesis that timberlines descend in altitude by around 1000 m during glacial cycles, the Sierra Madre Occidental likely harbored the largest, continuous extensions of timberline vegetation in Mexico during ice ages, ranging from central Sonora to southern Durango (McDonald 1993). In the present age, this refugium of highly specialized vegetation exhibits the narrowest distribution of all of Mexico's alpine habitats.

In anticipation of a comprehensive examination of the origins, floristic relationships, endemic composition, and historical biogeography of alpine vegetation throughout Mexico (McDonald & Feria, in prep), we present a floristic list for a rare and vulnerable plant community (by virtue of its narrow distribution) in the Sierra Madre Occidental. We compare this flora with a comparable vegetation type in Northeast Mexico (McDonald 1990) and that of Mexico's volcanic belt (Almeida et al. 2007), computing a 'Sorensen Index of Similarity' of only 7.2% between Cerro Mohinora and timberline vegetation of the Sierra Madre Oriental (SI = $2C/A + B$: where A and C equal the number of species found at disjunct sites and C equals the number of species found at both locations). In comparing the floras of Cerro Mohinora with a floristic list for Popocatepetl, Iztaccihuatl, and the Nevado de Toluca, the Sorensen Index computes to a mere 3.6% similarity. In both cases, floristic evidence indicates that the cyclic expansion of alpine vegetation during pluvial periods of the Pleistocene allows for only limited floristic exchange between the Sierra Madre Occidental and other tall mountains of Mexico. These data add further support to the general assumption that timberlines descend around 1000 m during glacial maxima.

Given the distinctive and rare character of Cerro Mohinora's plant communities, the summit and ecologically intact surroundings merit some form of conservation status. It is noteworthy that only a single non-native species (*Poa annua*) was encountered in the region during the late 1980s, indicating a relatively pristine flora, albeit affected by the grazing of goats during the 1950s (fide Correll) and cows during our two forays. A road now leads to the top of the mountain to serve a microwave tower, thereby encouraging and facilitating increased human activities in this isolated region. Foresight and planning might protect this rare, endemic, and predictably fragile vegetation type from fires, grazing, trail erosion, and other forms of ecological damage suffered by alpine zones throughout Mexico (Gimenez de Azcarate & Escamilla 1999; Almeida-Lenero et al. 2007).

ANNOTATED CHECKLIST OF VASCULAR PLANT TAXA

Collector names are abbreviated as follows: **DSC** = Donovan S. Correll (TEX); **GN** = Guy Nesom (TEX); **HSG** = Howard S. Gentry (TEX); **JM** = Jorge Martínez (HUAA, XAL); **JAM** = J. Andrew McDonald (HUAA, TEX, XAL). Herbarium acronyms denote location of each collector's specimens.

PTERIDOPHYTA

Woodsiaceae

Woodsia philipsii Windham; DSC & HSG 23164

CONIFEROPHYTA

Cupressaceae

Juniperus blancoi Marti var. *huehuetentensis* R.P. Adams, S. González & M. González Elizondo; JAM & JM 2351

Pinaceae

Pinus cooperi Blanco; JAM & JM 2349

MAGNOLIOPHYTA

Apiaceae

Eryngium lemmonii J.M. Coult. & Rose; JAM & JM 2364

Ligusticum porteri J.M. Coult. & Rose; JAM & JM 2404, JAM & GN 2471, GN & JAM 6458

Tauschia edulis (S. Wats.) J.M. Coult. & Rose; JAM & JM 2357

Brassicaceae

Pennellia longifolia (Benth.) Rollins; DSC & HSG 23167, JAM & JM 2353, JAM & GN 2480

Draba rubricaulis Heller; DSC & HSG 23160, JAM & JM 2405

Caryophyllaceae

Arenaria lanuginosa (Michx.) Rohrb.; DSC & HSG 23187, JAM & JM 2350, 2408, JAM & GN 2478

Drymaria effusa var. *depressa* (Greene) J. Duke; JAM & JM 2365

Cerastium nutans Raf.; JAM & JM 2366

Silene laciniata Cav. var. *greggii* (A. Gray) Hitch. & Maguire; DSC & HSG 23162, JAM & JM 2382, JAM & GN 2492

Commelinaceae

Commelina cf. *tuberosa* L.; JAM & JM 2377

Tradescantia pygmaea Hunt; JAM & JM 2378

Compositae

Ageratina lemmonii (B.L. Rob.) King & Rob.; JAM & JM 2374

Correllia montana A.M. Powell; JAM & JM 2346, JAM & GN 2479

Dahlia sherffii Sorenson; JAM & JM 2373

Erigeron macdonaldii G.L. Nesom; JAM & GN 2372

Erigeron mohinorensis G.L. Nesom; DSC & HSG 23151, JAM & JM 2371

Erigeron rhizomactis G.L. Nesom; JAM & JM 234X

Galinsoga subdiscoidea Cronquist; JAM & JM 2344, GN & JAM 6478

Grindelia megalcephala (Fern.) G.L. Nesom; *JAM & JM* 2347
Senecio mohinorensis Greenm.; *DSC & HSG* 23146, *JAM & JM* 2338
Senecio umbraculifera S. Wats.; *DSC & HSG* 23155, *JAM & JM* 2348
Senecio scalaris Greenm. var. *scalaris*; *DSC & HSG* 23144, *JAM & JM* 2342
Stevia plummerae A. Gray var. *plummerae*; *JAM & JM* 2339
Verbesina longifolia A. Gray; *JAM & JM* 2337
Viguiera multiflora S.F. Blake var. *macrocephala* (Heiser) B.L. Turner; *JAM & JM* 2375

Crassulaceae

Sedum jaliscanum S. Wats.; *JAM & GN* 2468
Sedum chihuahuense S. Wats.; *DSC & HSG* 23158, *JAM & JM* 2400, *JAM & GN* 2477
Villadia pringlei Rose; *JAM & JM* 2379T, *JAM & GN* 247

Cyperaceae

Cyperus pennellii O'Neill & Ben. Ayers; *JAM & JM* 2356 (XAL)
Cyperus fendlerianus Boeckeler; *JAM & JM* 2397 (XAL)

Fabaceae

Lupinus barberi (C.P.Smith) Rafaill & Dunn; *JAM & JM* 2354
Lupinus montanus Kunth subsp. *glabrior* (S. Watson) Dunn & Harmon; *JAM & JM* 2359

Gentianaceae

Halenia recurva (Sm.) Allen; *JAM & JM* 2396
Gentianopsis macrantha (D. Don ex G. Don) Iltis; *DSC & HSG* 23143

Geraniaceae

Geranium wislizeni S. Wats.; *DSC & HSG* 23159
Geranium richardsonii James; *JAM & JM* 2380

Hydrophyllaceae

Phacelia platycarpa Cav.; *JAM & JM* 2355

Iridaceae

Sisyrinchium convolutum Nocca; *JAM & JM* 2352

Juncaceae

Luzula comosa Meyer; *DSC & HSG* 23173, *JAM & JM* 2391, 2481, *JAM & GN* 2494

Lamiaceae

Salvia microphylla Kunth; *JAM & JM* 2385
Stachys mohinora B.L. Turner; *DSC & HSG* 23181, *JAM & JM* 2387, *JAM & GN* 2495

Liliaceae

Calochortus venustus Greene; *JAM & JM* 2369

Melanthaceae

Zigadenus cf. elegans Pursh; *JAM & JM* 2410

Montiaceae

Montia perfoliata (Donn ex Willd.) Howell; *JAM & JM* 2368, *JAM & GN* 2474

Onagraceae

Oenothera pubescens Willd. ex Spreng.; *JAM & JM* 2367

Orobanchaceae

Castilleja tenuifolia Mart. & Gal.; *DSC & HSG* 23147

Pedicularis chihuahuensis G.L. Nesom; *JAM & JM* 2402

Oxalidaceae

Oxalis decaphylla Kunth.; *JAM & JM* 2409

Piperaceae

Peperomia campyloptropa A.W. Hill; *JAM & JM* 2493

Plantaginaceae

Penstemon barbatus (Cav.) Roth.; *JAM & JM* 2386
Penstemon campanulatus (Cav.) Willd.; *DSC & HSG* 17996
Penstemon mohinoranus Straw; *JAM & JM* 2370
Veronica peregrina L. ssp. *xalapensis* (Kunth) Penn.; *JAM & JM* 2399

Poaceae

Bromus ciliatus L.; *JAM & JM* 2393, *GN & JM* 6474
Deschampsia flexuosa (L.) Trin.; *DSC & HSG* 23179, *JAM & GN* 2482
Festuca hintoniana Alexeev; *JAM & GN* 2475, 2489
Festuca diclina S.J. Darbyshire; *JAM & JM* 2392
Koeleria pyramidata (Lam.) Beauv.; *DSC & HSG* 23175, *JAM & JM* 2394
Poa annua L.; *JAM & JM* 2358
Trisetum filifolium Scribn. ex Beal; *DSC & HSG* 23176

Polygonaceae

Rumex acetosella L.; *JAM & JM* 2363

Primulaceae

Dodecatheon pulchellum var. *pulchellum* (Raf.) Merr.; *JAM & GN* 2490, *GN & JAM* 6460
Dodecatheon ellisiae Standl.; *GN & JAM* 6471
Primula rusbyi Greene; *JAM & JM* 2406, *GN & JAM* 6469

Ranunculaceae

Ranunculus gentryanus Benson; *DSC & HSG* 23156, *JAM & JM* 2362
Thalictrum grandifolium S. Wats.; *JAM & JM* 2383, *JAM & GN* 2470, *GN & JAM* 6457

Rosaceae

Alchemilla aphanoides L.f.; *JAM & JM* 2381
Holodiscus discolor (Pursh) Maxim. var. *discolor*; *DSC & HSG* 23182, *JAM & JM* 2403

Rubiaceae

Galium uncinulatum DC.; *JAM & GN* 2401, 2476
Galium mexicanum Kunth subsp. *asperimum* (A. Gray) Dempster; *JAM & JM* 2388

Salicaceae

Populus tremuloides Michx.; *DSC & HSG* 23186

Saxifragaceae

Heuchera sp. *JAM & JM* 2389
Saxifraga eriophora S. Wats.; *JAM & JM* 2407

Solanaceae

Solanum stolonifera Schtdl. & Bché; *JAM & JM* 2389, 2420
Solanum demissum Lindl.; *JAM & GN* 2469

Valerianaceae

Valeriana deltoidea Meyer; *JAM & JM* 2361

ACKNOWLEDGMENTS

Assistance in plant collection determinations were provided by Robert Adams, Rebecca van Devender, Peter Hoch, the late Charlotte and John Reeder, James Reveal, Gordon Tucker and Billie Turner. Tom Wendt (TEX) provided help on numerous occasions in accessing critical plant collections and floristic databases, while field vehicles, with financial and herbarium support for undertaking fieldwork, were provided the Universidad

Autónoma de Aguascalientes (HUAA) and the I.N.I.R.E.B. (now the Instituto de Ecología) in Xalapa, Veracruz (XAL). We thank Richard S. Felger and Alicia Melgoza for their helpful comments on the manuscript.

REFERENCES

- ALMEIDA-LENERO, L., A.M. CLEEF, A. HERRERA, A. VELAZQUEZ, AND I. LUNA. 1994. El zacatonal alpino del Volcán Popocatepetl, México, y su posición en las montañas tropicales de América. *Phytocoenologia* 22:391–436.
- ALMEIDA-LENERO, L., GIMENEZ DE AZCARATE, A.M. CLEEF, AND A. GONZALEZ TRAPAGA. 2004. Las comunidades vegetales del zacatonal alpino de los volcanes Popocatepetl y Nevado de Toluca, región central de México. *Phytocoenologia* 34:91–132.
- ALMEIDA-LENERO, L., M. ESCAMILLA, J. GIMENEZ DE AZCARATE, A. GONZALEZ TRAPAGA, AND A.M. CLEEF. 2007. Vegetación alpina de los volcanes Popocatepetl, Iztaccíhuatl y Nevado de Toluca. In: I. Luna, J.J. Morrone y D. Espinosa, eds. *Biodiversidad de la Faja Volcanica Transmexicana*. Universidad Nacional Autónoma de México, México, D.F. Pp. 179–198.
- BEAMAN, J. 1962. The timberlines of Iztaccíhuatl and Popocatepetl, Mexico. *Ecology* 42:377–385.
- BEAMAN, J. 1965. A preliminary ecological study of the alpine flora of Popocatepetl and Iztaccíhuatl. *Bol. Soc. Bot. Mexico* 29:63–75.
- BEAMAN, J. AND J.W. ANDRESEN. 1966. The vegetation, floristics and phytogeography of the summit of Cerro Potosi, Mexico. *Amer. Midl. Naturalist* 75:1–33.
- CORRELL, D.S. 1960. A mule-train trip to Sierra Mohinora, Chihuahua. *Amer. Fern J.* 50:66–78.
- GIMENEZ DE AZCARATE, J. AND M. ESCAMILLA. 1999. Las comunidades edafoferófilas (enebrales y zacatonales) en las montañas del centro de México. *Phytocoenologia* 29(4):449–468.
- GOLDMAN, E.A. 1951. Biological investigations in México 1892–1906. *Smithsonian Misc. Collect.* 115:127–129.
- LAUER, W. AND D. KLAUS. 1975. Geocological investigations on the timberline of Pico de Orizaba, Mexico. *Arctic Alpine Res.* 7:315–330.
- MCDONALD, J.A. 1990. The alpine-subalpine flora of northeastern Mexico. *Sida* 14:21–28.
- MCDONALD, J.A. 1993. Phytogeography and history of the alpine-subalpine flora of northeastern Mexico. In: T.P. Ramamoorthy, J. Fa, R. Bey, and A. Lot, eds. *Biological diversity of Mexico: origin and distribution*. Oxford University Press, New York. Pp. 681–703.
- NARAVE, H. 1985. La vegetación del Cofre de Perote, Veracruz, México. *Biótica* 1:35–57.
- PURPUS, C.A. 1907. Mexikanische hochgipfel. *Vegetationsbilder* 5:46–51.
- RZEDOWSKI, J. 1975. An ecological and phytogeographical analysis of the grasslands of Mexico. *Taxon* 24:67–80.
- WEBB, R.G. AND R.H. BAKER. 1984. Terrestrial vertebrates of the Cerro Mohinora Region, Chihuahua, Mexico. *Southw. Naturalist* 29:243–246.